

# Topical calcineurin inhibitors in dermatology.

## Part I: Properties, method and effectiveness of drug use

Katarzyna Gutfreund, Wojciech Bienias, Anna Szewczyk, Andrzej Kaszuba

Department of Dermatology, Pediatric Dermatology and Dermatological Oncology, Medical University of Lodz, Poland  
Head: Prof. Andrzej Kaszuba MD, PhD

Postep Derm Alergol 2013; XXX, 3: 165–169

DOI: 10.5114/pdia.2013.35619

### Abstract

Topical calcineurin inhibitors (TCI) are a relatively new class of drugs used in dermatology. There are two drug forms available – tacrolimus 0.03% or 0.1% ointment and 1.0% pimecrolimus cream. The drugs act by inhibiting synthesis of proinflammatory cytokines. The only approved indication for using TCI is treatment of atopic dermatitis. The TCI may be used as an alternative therapy to corticosteroids. Tacrolimus is used to treat moderate-to-severe atopic dermatitis, pimecrolimus – mild-to-moderate atopic dermatitis. Topical calcineurin inhibitors do not cause skin atrophy and the drug absorption through the skin is minimal. The TCI have been well-studied, their efficacy was evaluated in a number of vast, long-term studies. The anti-inflammatory potency of tacrolimus ointment is similar to a corticosteroid with moderate activity, while the latter is clearly more active than pimecrolimus cream. Topical calcineurin inhibitors significantly relieve pruritus in atopic eczema.

**Key words:** topical calcineurin inhibitors, tacrolimus, pimecrolimus.

### Introduction

The history of calcineurin inhibitors is associated with search for immunosuppressive drugs for transplantation. Cyclosporine A was the first introduced calcineurin. Transplants of kidneys, liver, lungs or heart were possible due to drug inhibition of immune reaction against the transplanted organ. Improvement in clinical presentation of skin was observed in patients with psoriasis, atopic dermatitis and other dermatoses when given cyclosporine A after the organ transplant. Cyclosporine A is still used in treating some severe skin conditions but in transplantation it was replaced by tacrolimus, isolated in 1984, for its 10-100 times higher immunosuppressive activity [1]. Tacrolimus is produced by a type of soil bacterium *Streptomyces tsukubaensis* in the *Tsukuba* region of Japan. First known as FK 506, then the name tacrolimus was derived from *Tsukuba* (the mountain the soil sample came from), *macrolide* (the compound is a hydrophobic macrolide) and *immunosuppressive* [1, 2]. Tacrolimus was introduced as an oral drug preventing transplant rejection in 1989. Despite several attempts to treat atopic dermatitis, psoriasis, alopecia areata, pemphigus or eosinophil fasciitis with the drug, it is not commonly used in systemic treatment in derma-

tology, which may be due to its high costs [3]. Discovery of pimecrolimus, another calcineurin inhibitor, resulted from long studies on ascomycin derivatives (antifungal and immunomodulatory compound) in the Laboratory of Novartis. Pimecrolimus, ascomycin macrolactam, is produced by the fermentation of *Streptomyces hygroscopicus* var. *ascomycetous*.

Tacrolimus (topical) was launched on the market at the end of 2000, pimecrolimus in 2001. Attempts of topical use of cyclosporine A were not successful due to high molecular weight (about 1200 Da), preventing effective penetration into the skin [4]. Sporadically, the drug is used in the form of eye drops in eye inflammation and for washing the oral cavity in patients with pemphigus vulgaris [5, 6]. Tacrolimus and pimecrolimus have similar both chemical structure and molecular mass (about 800 Da). The molecular structure allows effective skin penetration, not as intense as topical glucocorticosteroids, which prevents higher systemic drug absorption [7].

### Mechanism of action of calcineurin inhibitors

Mechanism of action of calcineurin inhibitors is suppressing synthesis of pro-inflammatory cytokines. In the

---

**Address for correspondence:** Prof. Andrzej Kaszuba MD, PhD, Department of Dermatology, Pediatric Dermatology and Dermatological Oncology, Medical University of Lodz, 1/5 Kniaziewiczza St, 91-347 Lodz, Poland, phone: +48 42 651 10 72, e-mail: andrzej.kaszuba@umed.lodz.pl  
**Received:** 15.03.2013, **accepted:** 4.06.2013.

cytoplasm of the target cells, pimecrolimus and tacrolimus bind to the intracellular protein macrophilin-12, also called FKBP (FK506-binding protein). Similarly, cyclosporine A binds to cyclophilin. Immunosuppressive activity results from suppressing calcineurin activity – dependent on calcium and calmodulin (serine-threonine phosphatase). The drug has an anti-inflammatory activity due to T-helper activity affecting synthesis and release of pro-inflammatory cytokines. Cytokine transcription blockage leads to decrease in expression of cytokine Th1 and Th2 dependent among others on interleukin 2, 3, 4 and 5 (IL-2, IL-3, IL-4, IL-5), interferon- $\gamma$  (INF- $\gamma$ ) and tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ). Tacrolimus and pimecrolimus inhibit mast cell and neutrophil activation and release of inflammatory mediators. Tacrolimus affects basophils and eosinophils function as well as function and apoptosis induction of Langerhans cells [7, 8].

### Indications for use of topical calcineurin inhibitors

Tacrolimus is available in the form of 0.1% and 0.03% ointment (Protopic) while pimecrolimus in the form of 1% cream (Elidel). The only approved indication for use of topical calcineurin inhibitors (TCI) is atopic dermatitis (AD). However, these preparations are also very effective in treating other dermatoses, particularly seborrheic dermatitis, genital lichen sclerosus, oral lichen planus, psoriasis (face and flexures areas), vitiligo and alopecia areata [9, 10].

Tacrolimus is used for treating moderate-to-severe AD when there is no sufficient response or tolerance to conventional treatment, such as topical use of corticosteroids. The drug was also approved for supportive therapy (proactive therapy) of moderate-to-severe AD to prevent relapses and extend periods without recurrence in patients with frequent exacerbations (4 or more times a year) and who initially used to respond to treatment with tacrolimus ointment 2 times a day for maximum 6 weeks.

The new proactive therapy has been based on new discoveries in AD pathogenesis. In the course of this inflammatory dermatosis, disorders affecting both the structure and immune function are observed in seemingly unchanged skin. They are reported to be significantly dependent on filaggrin gene mutation. After disappearance of active lesions, the skin still presents features of subclinical inflammation [11].

In remission tacrolimus should be applied on the skin area where most intense lesions were observed in the course of the last exacerbation or where lesions usually appear, one or two times a day for 2 days a week (e.g. Monday and Thursday). Significant effectiveness of proactive therapy in preventing AD exacerbations and lower cost compared with conventional reactive treatment are stressed [12]. So far proactive therapies for the described tacrolimus as well as for two glucocorticosteroids: fluticasone and methylprednisolone have been investigated [11].

Pimecrolimus was approved for mild-to-moderate AD when topical corticosteroids are contraindicated or not tolerated. Pimecrolimus in proactive therapy has not been studied.

Both topical calcineurin inhibitors were approved for 2+ years. Pimecrolimus should be applied on lesions two times a day until they subside, but not longer than 6 weeks. Tacrolimus (0.03%) is indicated for patients between 2 and 16 years and is applied 2 times a day for 3 weeks and then one time a day until disappearance of symptoms. Tacrolimus (0.1%) is for 16+ years, applied 2 times a day until disappearance of symptoms (Table 1).

### Glucocorticosteroids and topical calcineurin inhibitors

Topical calcineurin inhibitors should be used as a second-line drugs following topical glucocorticosteroids (GCs), which are regarded as topical treatment “gold standard” for the disease exacerbations [13]. The great advan-

**Table 1.** The use of tacrolimus and pimecrolimus

Tacrolimus	Pimecrolimus
<ul style="list-style-type: none"> <li>Treatment of <b>moderate-to-severe</b> AD in case there is no satisfactory response or no tolerance to conventional treatment, such as topical corticosteroids</li> <li>From 2 to 16 years, first 0.03% ointment twice daily for 3 weeks, then once daily until disappearance of lesions</li> <li>From 16 years, 0.1% ointment until disappearance of lesions</li> <li><b>Supportive treatment of moderate-to-severe</b> AD to prevent recurrence and extend periods without recurrence in patients with frequent exacerbations (4 or more times a year) who initially responded to treatment with tacrolimus ointment twice daily for max. 6 weeks</li> <li>Use once daily 2 days a week (e.g. Monday and Thursday) on the affected surface</li> <li>0.1% ointment in adults, 0.03% in children</li> </ul>	<ul style="list-style-type: none"> <li><b>Mild-to-moderate</b> AD when topical corticosteroids are not recommended or not possible</li> <li>Adults and children older than 2 years, twice daily until lesions disappear, not longer than 6 weeks</li> </ul>

**Table 2.** Mechanism of action of pimecrolimus, tacrolimus and topical corticosteroids

Parameter	Pimecrolimus	Tacrolimus	Corticosteroids
Activity on cells	T lymphocytes Mast cells	T lymphocytes Mast cells Eosinophils, basophils Langerhans cells	T lymphocytes Mast cells Eosinophils, basophils Langerhans cells Keratinocytes Endothelial cells Fibroblasts
Cytokines	IL-2, IL-3, IL-4, IL-5, INF- $\gamma$ , TNF- $\alpha$	IL-2, IL-3, IL-4, IL-5, INF- $\gamma$ , TNF- $\alpha$	IL-2, IL-3, IL-4, IL-5, INF- $\gamma$ , TNF- $\alpha$ , INF- $\alpha$ , GM-CSF
Function of blocking and apoptosis of Langerhans cells	–	+	++
Absorption through the skin	+	++	+++
Atrophogenic activity	–	–	+++

tage of topical glucocorticosteroids is their fast and effective action, however, there are adverse effects to be considered. Long-term use of GCs may lead to skin atrophy, telangiectasia, striae, hyper- and hypopigmentation, as well as to anaphylaxis and contact allergy. In extreme cases, excessive overuse of topical GCs may cause hypothalamic-pituitary-adrenal suppression, induce Cushing's syndrome. There were cases of developing glaucoma in children following GCs application on the lid [14]. In contrast, TCI due to their high affinity for the receptor and lower absorption through skin do not cause such symptoms. The TCI are especially indicated in long-term therapies or areas such as the face, folds and genital region, where penetration through skin is many times higher [13].

The TCI do not have atrophogenic activity as they affect neither endothelial cells nor fibroblasts like topical GCs. Target cells for pimecrolimus are T lymphocytes, mast cells, whereas for tacrolimus – T lymphocytes, mast cells, eosinophils, basophils and most importantly Langerhans cells [7]. Kyllönen *et al.* found that long-term therapy with tacrolimus leads to withdrawal of steroid-dependent skin atrophy and increase in collagen synthesis and skin thickness [15] (Table 2).

### Pharmacokinetics of topical calcineurin inhibitors

The TCI absorption through the skin into circulation is minimal due to a large molecular size of the drugs. The highest absorption is observed in the initial stage of AD treatment when the inflammatory process is most advanced. When the disease subsides and epidermal barrier recovers, penetration into the skin decreases [16]. Increased tacrolimus penetration into circulation was found in dermatoses where profound skin barrier damage is observed e.g. Netherton syndrome [17], lamellar ichthyosis [18], pyoderma gangrenosum [19]. Pimecrolimus

is more lipophilic than tacrolimus, which results in slow penetration of TCI from the corneal layer rich in lipids into the hydrated lower epidermal layer [20]. The TCI penetrates much lower than topical glucocorticosteroids [21].

Draeos *et al.* compared pimecrolimus and tacrolimus concentrations in the blood from patients given one of the preparations 2 times a day for 13 days. Tacrolimus was detectable in 36% of blood samples whereas pimecrolimus in 12% and the concentration was very low. Subsequent studies on TCI pharmacokinetics confirmed very low absorption into systemic circulation [22]. Undre *et al.* evaluated tacrolimus distribution in the skin and serum after application on the extended surface of the skin. Gradual decrease in the drug concentration in the skin along with the increasing skin depth was found, moreover, the retention time was short after the drug withdrawal. Systemic drug activity was not confirmed; in 64% of patients the concentration was below 1 ng/ml, and in 30% it was undetectable [23]. Pharmacokinetic studies on pimecrolimus carried out on children with moderate-to-severe AD and whose body surface was affected up to 92%, showed 67% of pimecrolimus blood concentrations below 0.5 ng/ml and 97% of samples did not exceed 2.0 ng/ml [24]. In a recent study carried out on a group of very young children (between 3 and 24 months) given 0.03% tacrolimus 2 times daily for 2 years in exacerbation, the concentration of tacrolimus was < 1.0 ng/ml in 98% of blood samples and in > 40% of blood samples, tacrolimus concentration was below the limit of detection (0.0250 ng/ml) [25].

In the investigated groups of children there was no difference in drug absorption through skin depending on age. No increased risk of systemic or local infection was found. Proper immune response to vaccination in the course of treatment with TCI was observed, which confirms only topical activity of the drug [26].

### Efficacy of topical calcineurin inhibitors

The TCI are a well-investigated class of drugs, their efficacy was assessed in many longitudinal studies on vast populations of patients. First clinical reports on topical tacrolimus and pimecrolimus described the drug significant efficacy in the management of patients with AD.

In the study on a group of 624 children with moderate-to-severe AD, Reitamo *et al.* found that 0.03% tacrolimus ointment is more effective than 1% hydrocortisone acetate [27]. The following investigations carried out by Remitz *et al.* on a group of 466 children at the age of 2-15 years for 30 months showed effectiveness of tacrolimus in AD management [28]. In 2008, results from a multi-centre study on the effectiveness and safety of tacrolimus use, conducted on a group of 782 patients with AD at the age between 2 and 72 years, were published. The treatment was stopped when all skin lesions and pruritus disappeared, in the case of recurrence, the therapy was restarted. After 4 weeks' therapy, 60% of patients or their caretakers evaluated the clinical picture as very good or excellent. In the final stage 75% of participants of the study were satisfied with the treatment [29].

The obtained results from the studies on effectiveness of pimecrolimus are also satisfying. In a double-blind study on a group of 713 children between 2 and 17 years with AD carried out for 12 months, Wahn *et al.* compared effectiveness of treatment with pimecrolimus to conventional topical methods (glucocorticosteroids, emollients). Clinical evaluation was performed after 6 and 12 months of treatment, safety and tolerance to the preparations were also monitored; improvement in the clinical picture was found after both 6 and 12 months of the treatment [30]. In the comparative study of pimecrolimus and the drug vehicle, Kaufmann *et al.* proved significant reduction in pruritus; improvement was observed in 56% of patients applying pimecrolimus and 34% of patients given placebo [31]. Reduction in pruritus after using 1% pimecrolimus was also found in a study on 186 children with mild-to-moderate AD conducted by Ho *et al.* [32]. In the study on adult patients with AD, Meurer *et al.* found decrease in Eczema Area and Severity Index (EASI) by 33% in the first 7 days and by 69.8% in the 6<sup>th</sup> week of treatment with pimecrolimus. In contrast, there was an increase in EASI by 3.8% in the first week and by 15.9% in the 6<sup>th</sup> week in the control group using drug vehicle [33].

Paller *et al.* conducted comparative studies on effectiveness of tacrolimus and pimecrolimus in children and adult patients with AD. Randomized single-blinded studies were carried out on 1065 patients for 6 weeks by three multi-study centres. The obtained results suggest that tacrolimus was significantly more effective than pimecrolimus in EASI both in children and adult patients with moderate-to-severe AD. Tacrolimus action started earlier compared with pimecrolimus [34].

Studies indicate that anti-inflammatory activity of tacrolimus may be compared with activity of medium poten-

cy corticosteroids while activity of pimecrolimus is less potent than that of tacrolimus. Both TCI are effective in pruritus reduction, which is a great advantage in treating patients with AD suffering from this symptom [13].

### References

1. Ruzicka T, Assmann T, Homey B. Tacrolimus: the drug for the turn of the millennium? *Arch Dermatol* 1999; 135: 574-80.
2. Gupta AK, Adamiak A, Chow M. Tacrolimus: a review of its use for the management of dermatoses. *J Eur Acad Dermatol Venereol* 2002; 16: 100-14.
3. Tharp MD. Calcineurin inhibitors. *Dermatol Ther* 2002; 15: 325-32.
4. Surber C, Itin P, Büchner S. Clinical controversy on the effect of topical cyclosporin: what is the target site? *Dermatology* 1992; 185: 242-5.
5. Brown MM, Brown GC, Brown HC, et al. Value-based medicine, comparative effectiveness, and cost-effectiveness analysis of topical cyclosporine for the treatment of dry eye syndrome. *Arch Ophthalmol* 2009; 127: 146-52.
6. Pacor ML, Biesi D, Carleto A, et al. Topical cyclosporine in the treatment of oral pemphigus. *Minerva Stomatol* 1998; 47: 183-6.
7. Nighiem P, Pearson G, Langley RG. Tacrolimus and pimecrolimus: from clever prokaryotes to inhibiting calcineurin and treating atopic dermatitis. *J Am Acad Dermatol* 2002; 46: 228-41.
8. Gisondi P, Ellis CN, Girolomoni G. Pimecrolimus in dermatology: atopic dermatitis and beyond. *Int J Clin Pract* 2005; 59: 969-74.
9. Wollina U. The role of topical calcineurin inhibitors for skin diseases other than atopic dermatitis. *Am J Clin Dermatol* 2007; 8: 157-73.
10. Silny W, Sadowska A, Dańczak-Pazdrowska A, Polańska A. Application of tacrolimus in the treatment of skin diseases other than atopic dermatitis. *Postep Derm Alergol* 2011; 28: 41-5.
11. Wollenberg A, Bieber T. Proactive therapy of atopic dermatitis – an emerging concept. *Allergy* 2009; 64: 276-8.
12. Healy E, Bentley A, Fidler C, et al. Cost-effectiveness of tacrolimus ointment in adults and children with moderate and severe atopic dermatitis: twice-weekly maintenance treatment vs. standard twice-daily reactive treatment of exacerbations from a third party payer (U.K. National Health Service) perspective. *Br J Dermatol* 2011; 164: 387-95.
13. Ring J, Alomar A, Bieber T, et al. Guidelines for treatment of atopic eczema (atopic dermatitis) part I. *J Eur Acad Dermatol Venereol* 2012; 26: 1045-60.
14. Garrott HM, Walland MJ. Glaucoma from topical corticosteroids to the eyelids. *Clin Exp Ophthalmol* 2004; 32: 224-6.
15. Kyllönen H, Remitz A, Mandelin JM, et al. Effects of 1-year intermittent treatments with topical tacrolimus monotherapy on skin collagen synthesis in patients with atopic dermatitis. *Br J Dermatol* 2004; 150: 1174-81.
16. Rustin MH. The safety of tacrolimus ointment for the treatment of atopic dermatitis: a review. *Br J Dermatol* 2007; 157: 861-73.
17. Allen A, Siegfried E, Silverman R, et al. Significant absorption of topical tacrolimus in 3 patients with Netherton syndrome. *Arch Dermatol* 2001; 137: 747-50.

18. Allen DM, Esterly NB. Significant systemic absorption of tacrolimus after topical application in a patient with lamellar ichthyosis. *Arch Dermatol* 2002; 138: 1259-60.
19. Pitarch G, Torrijos A, Mahquijes L, et al. Systemic absorption of topical tacrolimus in pyoderma gangrenosum. *Acta Derm Venereol* 2006; 86: 64-5.
20. Billich A, Aschauer H, Aszodi A, et al. Percutaneous absorption of drugs used in atopic eczema: pimecrolimus permeates less through skin than corticosteroids and tacrolimus. *Int J Pharm* 2004; 269: 29-35.
21. Weiss HM, Fresneau M, Moenius T, et al. Binding of pimecrolimus and tacrolimus to skin and plasma proteins: implications for systemic exposure after topical application. *Drug Metab Dispos* 2008; 36: 1812-8.
22. Draelos Z, Nayak A, Pariser D, et al. Pharmacokinetics of topical calcineurin inhibitors in adult atopic dermatitis: a randomized, investigator-blind comparison. *J Am Acad Dermatol* 2005; 53: 602-9.
23. Undre NA, Moloney FJ, Ahmadi S, et al. Skin and systemic pharmacokinetics of tacrolimus following topical application of tacrolimus ointment in adult with moderate to severe atopic dermatitis. *Br J Dermatol* 2009; 160: 665-9.
24. Hultsch T, Kapp A, Spergel J. Immunomodulation and safety of topical calcineurin inhibitors for the treatment of atopic dermatitis. *Dermatology* 2005; 211: 174-87.
25. Mandelin JM, Rubins A, Remitz A, et al. Long-term efficacy and tolerability of tacrolimus 0.03% ointment in infants: a two-year open-label study. *Int J Dermatol* 2012; 51: 104-10.
26. Hofman T, Cranswick N, Kuna P, et al. Tacrolimus ointment does not affect the immediate response to vaccination, the generation of immune memory, or humoral and cell-mediated immunity in children. *Arch Dis Child* 2006; 91: 905-10.
27. Reitamo S, Harper J, Bos JD, et al. 0.03% Tacrolimus ointment applied once or twice daily is more efficacious than 1% hydrocortisone acetate in children with moderate to severe atopic dermatitis: results of a randomized double-blind controlled trial. *Br J Dermatol* 2004; 150: 554-62.
28. Remitz A, Harper J, Rustin M, et al. Long-term safety and efficacy of tacrolimus ointment for the treatment of atopic dermatitis in children. *Acta Derm Venereol* 2007; 87: 54-61.
29. Reitamo S, Rustin M, Harper J, et al. 0.1% Tacrolimus Ointment Long-term Follow-up Study Group. A 4 year follow-up study of atopic dermatitis therapy with 0.1% tacrolimus ointment in children and adult patients. *Br J Dermatol* 2008; 159: 942-51.
30. Wahn U, Bos JD, Goodfield M, et al. Efficacy and safety of pimecrolimus cream in the long-term management of atopic dermatitis in children. *Pediatrics* 2002; 110: e2.
31. Kaufmann R, Bieber T, Helgesen AL, et al. Onset of pruritus relief with pimecrolimus cream 1% in adult patients with atopic dermatitis: a randomized trial. *Allergy* 2006; 61: 375-81.
32. Ho VC, Gupta A, Kaufmann R, et al. Safety and efficacy of nonsteroid pimecrolimus cream 1% in the treatment of atopic dermatitis in infants. *J Pediatr* 2003; 142: 155-62.
33. Meurer M, Fartasch M, Albrecht G, et al. Long-term efficacy and safety of pimecrolimus cream 1% in adults with moderate atopic dermatitis. *Dermatology* 2004; 208: 365-72.
34. Paller AS, Lebwohl M, Fleischer AB, et al. Tacrolimus ointment is more effective than pimecrolimus cream with a similar safety profile in the treatment of atopic dermatitis: results from 3 randomized, comparative studies. *J Am Acad Dermatol* 2005; 52: 810-22.